







Diphenhydramine (Antihistamine) Dosage Table (i.e., Benadryl®) allergic reactions, nasal allergies, and hives.

- May be given every 4 - 6 hours as directed by the doctor.
- Do NOT exceed more than 6 doses in 24 hours.
- children under 2 years: do not use
- children 2 to 5 years: do not use unless directed by a doctor

mL = milliliter
tsp = teaspoonful
mg = Milligram

Liquid Diphenhydramine for Children 12.5mg (in each 5mL or 1 tsp)

WEIGHT (Pounds)	DOSE Liquid for Children
20-24 lbs	4 mL (3/4 tsp) 
25-37 lbs	5 mL (1 tsp) 
38-49 lbs	7.5 mL (1 1/2 tsp) 
50-99 lbs	10 mL (2 tsp) 



Chewable Tablets for Children (12.5mg per tablet)

Weight (Pounds)	DOSE Number of Tabs for Children
Under 25 lbs	ASK THE DOCTOR
25-37 lbs	1
38-49 lbs	1 1/2
50-99 lbs	2
100 lbs +	4

For More Dosage Information, Including Tablets, Click on The Link Below

<https://bit.ly/2MYAFq3>

or

<https://www.healthychildren.org/English/safety-prevention/at-home/medication-safety/Pages/Diphenhydramine-Benadryl-Antihistamine.aspx>