





## Ibuprofen Dosage Table for Fever and Pain (i.e., Motrin®, Advil®)

- May be given every 6 hours.
- Do NOT exceed more than 4 doses in 24 hours.
- **We do not recommend Ibuprofen for children under 6 months of age.**
- If possible, use weight to dose; otherwise use age. Always ask a healthcare provider which product is right for your child.






mL = milliliter  
tsp = teaspoonful  
mg = Milligram

### Infant Concentrated Drops (50 mg/1.25mL)

WEIGHT (Pounds)	AGE (Estimated)	DOSE Infant Concentrated Drops
0-12 lbs	0-6 months	<b>DO NOT USE</b>
13-17 lbs	6-11 months	1.25 mL 
18-23 lbs	12-23 months	1.875 mL 



### Liquid for Children (100mg per 5mL or 1 tsp)

Weight (Pounds)	Age (Estimated)	DOSE Liquid for Children
0-12 lbs	0-6 months	DO NOT USE
13-17 lbs	6-11 months	2.5 mL (1/2 tsp)
18-23 lbs	12-23 months	4 mL (3/4 tsp)
24-35 lbs	2-3 years	5 mL (1 tsp) 
36-47 lbs	4-5 years	7.5 mL (1 1/2 tsp) 
48-59 lbs	6-8 years	10 mL (2 tsp) 
60-71 lbs	9-10 years	12.5 mL (2 1/2 tsp) 
72-95 lbs	11 years	15 mL (3 tsp) 

For More Dosage Information, Including Tablets, Click on The Link Below

<https://tinyurl.com/y565n67h>

or

<https://www.healthychildren.org/English/safety-prevention/at-home/medication-safety/Pages/lbuprofen-for-Fever-and-Pain.aspx>